(1)

(2) (3)

J. DEPA I MENI OF I HE INI E IO

J. . GEOLOGICAL J VEY

CLAGGIFICATION OF UATEENAEY GUEFICIAL DE OGITG, GOUTHEEN CALIFOENIA AEEAL MA ING EOJECT (GCAM) Alluvial Deposits

Version 1.0 09/10/2000

ALLJ VIAL DEPO II



VEEY YOUNG.--J ndissected surfaces; rilled to incised to veneered; older elements may have organic layer and (or) slight A horizon

YOUNG.--J ndissected to slightly dissected surfaces; A/Cox and Bcambic soils; slight to moderate K horizon; slight pavement & varnish; slight calcrete; ; in arid settings, Av horizon developed

OLD.--Moderately dissected surfaces; good Bt horizon; 10Y to 5Y hues; slightly sticky and plastic; continuous (±) clay films; moderate K horizon; moderate pavement and varnish; moderate calcrete

VEEY OLD.--Well dissected surfaces; strong Bt horizon with 5Y to 10 hues; sticky & plastic; prismatic structure; thick clay films on ped faces; strong K horizons; excellent pavement; good calcrete & duripan

Age Wash Alluvial-fan Alluvial-vallev lope-wash Pediment-veneer deposits deposits deposits deposits deposits Q4b Qw Qf Qa Qw_2 Qf2 Qaa Qsw Qpv 19 Q4a Qw₁ Qf₁ Qa₁ Holocene Qyf₅ Qva₅ 18 Qya₄ Qyf₄ Qyw Qyf Qyf₃ Qya Qya₃ Qysw Qypv 17 Qyf₂ Qya₂ 16 Qyf₁ Qya₁ 15 Qof₃ Qoa₃ 14 Qow Qof Qof₂ Qoa Qoa2 Qosw 13 Qopv Pleistocene Qoa₁ Qof₁ 12 Qvoa₃ Qvof₃ Qvoa₂ Qvopv 11 Qvow Qvof Qvof₂ Qvoa Qvosw Qvof₁ Qvoa₁

For CAMP geologic-map products, grain-size and physical-property information for surficial units is stored in digital data bases through the use of coded attributes (Matti and others, 1997). At the option of the geologic-map author, characteristic grain size information can be displayed in plot files through the use of alpha characters (e.g. Qyf_b, Qoa_{2a}), where the characters conform to the following definitions:

- a arenaceous (very coarse sand through very fine sand)
- b boulder gravel (25mm)
- g gravel (cobble through granule gravel)
- s siltv
- c clavev
- m- marl
- p peat

- (1) Numerical time scale is not linear;
- (2)I errace-age designations proposed by McFadden (1982) and by Bull (1991, Figure 4.11) for alluvial deposits in Mediterranean-climate regimes of southern California;
- (3) Geomorphic-surface designations proposed by Bull (1991, I able 2-13) in arid climatic regimes of southern California

Bull, W. ., 1991, Geomorphic responses to climatic change: New York, Oxford J niversity Press, 326 p. McFadden, L.D., 1982, I he impacts of temporal and spatial climatic changes on alluvial soils genesis in southern

Matti, J.C., Miller, F.K., Powell, .E., Kennedy, .A., and Ph.D. thesis, 430 p. Cossette, P.M., 1997a, Geologic-polygon attributes for digital geologic-map data bases produced by the outhern California Areal Mapping Project, version 1.0: J . . Geological urvey Open-File eport 97-860, 248 p.

McFadden, L.D., 1982, I he impacts of temporal and spatial climatic changes on alluvial soils genesis in southern California: I ucson, J niversity of Arizona, unpublished Ph.D. thesis, 430 p.